

Shading

Reading

Required

- Foley, Section 16.1

Optional

- Hearn & Baker, sections 14.1, 14.2, 14.5

Introduction

- So far, we've talked exclusively about geometry.
 - What is the shape of an object?
 - How do I place it in a virtual 3D space?
 - How do I know which pixels it covers?
 - How do I know which of the pixels I should actually draw?
- Once we've answered all those, we have to ask one more important question:
 - What value do I set each pixel to?
- Answering this question is the job of the **shading model**.
- (Of course, people also call it a lighting model, a light reflection model, a local illumination model, a reflectance model, etc., etc.)

Tedious Reality


- Properly determining the right color is *really hard*.
- Look around the room. Each light source has different characteristics. Trillions of photons are pouring out every second.
- These photons can:
 - interact with the atmosphere, or with things in the atmosphere
 - strike a surface and
 - be absorbed
 - be reflected
 - cause fluorescence or phosphorescence
- of course, none of the surfaces in here are perfect spheres or cylinders. At some microscopic level (very important for photons) they're all really bumpy.
- also, everything depends on wavelength.

Our Problem

- We're going to build up to an *approximation* of reality called the **Phong illumination model**.
- It has the following characteristics:
 - *not* physically based
 - gives a first-order *approximation* to physical light reflection
 - very fast
 - widely used


Ambient Light

R 0 G 0 B 0



Light Color

R 100 G 200 B 100




Object Color

R 150 G 50 B 50




Diffuse Color

R 0 G 0 B 0



Specular Color

R 0 G 0 B 0




Redraw

N 10 D 10 T 60



Ambient Light




Light Color




Object Color



Diffuse Color




Specular Color



Redraw





Ambient Light

R 250 G 150 B 150

R 100 G 200 B 100

Light Color

Object Color

R 150 G 50 B 50

R 160 G 155 B 155

Diffuse Color

R 100 G 200 B 100

Specular Color

R 0 G 0 B 0


Redraw

N 10 D 10 T 60

Ambient Light

R: 250 | G: 150 | B: 150

R: 100 | G: 200 | B: 100



Object Color

R: 150 | G: 50 | B: 50

Diffuse Color

R: 160 | G: 155 | B: 155

Specular Color

R: 204 | G: 198 | B: 198

Refraction

N: 10 | D: 10 | T: 60

Redraw

Ambient Light

R	250
G	150
B	150

Light Color

R	100
G	200
B	100

Object Color

R	150
G	50
B	50

Diffuse Color


R	179
G	171
B	130

Specular Color

R	204
G	198
B	198

Redraw

N	179
D	10
T	60



Iteration Zero


- Given:
 - a point P on a surface (P is determined by ray-object intersection, for instance)
 - visible through pixel p
- Assign each polygon a single color:

$$I = k_e$$

where

- I is the resulting intensity
 - k_e is the intrinsic shade associated with the object
- This has some special-purpose uses, but not really good for drawing a scene.

What will it like?



Iteration One


- Let's make the color at least dependent on the overall quantity of light available in the scene:

$$I = k_a I_a$$

- k_a is the **ambient reflection coefficient**.
 - really the reflectance of ambient light
 - “ambient” light is assumed to be equal in all directions
- I_a is the **ambient intensity**.
- Physically, what is “ambient” light?
 - Answer on next page.

Indirect Illumination (Ambient)

- Some surfaces are illuminated even it is in shadow. Why?
- There is indirect lighting (background lighting) reflected from other surfaces



- Each surface illuminated becomes itself a source of light for illumination on other surfaces of the scene.
- Each of these surfaces, in turn, reflect light to other surfaces, including the original one, thus achieving an “infinite regression” of reflections and illumination.
- Heuristic: Simply assume the indirect lighting is constant (same to all objects in the scene) in most graphics systems.

Wavelength Dependence

- Really, k_a and I_a are functions over all wavelengths κ .
- Ideally, we would do the calculation on these functions:

$$I(\kappa) = k_a(\kappa) I_a(\kappa)$$

- then we would find good RGB values to represent the spectrum $I_a(\kappa)$.
- Traditionally, though, k_a and I_a are represented as RGB triples, and the computation is performed on each color channel separately.

Diffuse Reflection

- Let's examine the ambient shading model:
 - objects have different colors
 - we can control the overall light intensity
 - what happens when we turn off the lights?
 - what happens as the light intensity increases?
 - what happens if we change the color of the lights?
- So far, objects are uniformly lit.
 - not the way things really appear
 - in reality, light sources are directional
- Diffuse, or **Lambertian** reflection will allow reflected intensity to vary with the direction of the light.

Diffuse Reflector


- Diffuse reflection occurs from dull, matte surfaces, like latex paint, or chalk.
- These **diffuse** or **Lambertian** reflectors reradiate light equally in all directions.
- Picture a rough surface with lots of tiny **microfacets**.



- Note:
 - Light may actually penetrate the surface, bounce around, and then reflect back out.
 - Accounts for colorization of diffusely reflected light by plastics.

Diffuse Reflector

- The reflected intensity from a diffuse surface does not depend on the direction of the viewer. The incoming light, though, does depend on the direction of the light source.



- Q: Why is the North Pole cold? Why is winter cold?

Iteration Two

- The incoming energy is proportional to $\cos \eta$, giving the diffuse reflection equations:

$$\begin{aligned} I &= k_e + k_a I_a + k_d I_l \cos \eta \\ &= k_e + k_a I_a + k_d I_l (\mathbf{N} \cdot \mathbf{L})_+ \end{aligned}$$

- where:
 - k_d is the **diffuse reflection coefficient**.
 - I_l is the intensity of the light source
 - \mathbf{N} is the normal to the surface (unit vector)
 - \mathbf{L} is the direction to the light source (unit vector)
 - $(x)_+$ means $\max\{0, x\}$
- OpenGL supports different kinds of lights: point, directional, and spot. How do these work?

Ideal Light Sources

Light Sources:

- In computer graphics, two types of light sources are commonly used
- point source The light source is a zero-volume point
- directional source The point source that are infinite far away




- Both types of light sources are ideal light sources (i.e. not realistic)
- But they are easy for computation.

Extended Light Sources


- A more realistic class of light source is **extended source**.
- The light source is actually a surface or a volume, not a point.

e.g

Fluorescent Lamp




Sun




- Sun is really an extended source
- Fluorescent light is a typical extended source.

Ambient and Diffuse Examples

- Increasing the diffuse coefficient:



- Increasing the ambient term while keeping the diffuse term constant:



Intensity drop-off with distance

- The laws of physics state that the intensity of a point light source must drop off with its distance squared.
- We can incorporate this effect by multiplying I_1 by $1/d^2$.
- Sometimes, this distance-squared drop off is considered too “harsh.” Angel suggests using with user-supplied constants for a , b , and c .

$$f(d) = \frac{1}{a + bd + cd^2}$$

- with user-supplied constants for a , b , and c .


$$f(d) = \min \left(1, \frac{1}{a + bd + cd^2} \right)$$

Specular Reflection

- **Specular reflection** accounts for the highlight that you see on some objects.
- It is particularly important for *smooth, shiny* surfaces, such as:
 - metal
 - polished stone
 - plastics
 - apples
- Specular reflection depends on the viewing direction V . The color is often determined solely by the color of the light.
 - corresponds to absence of internal reflections


Specular Reflection Derivation

- For a perfect mirror reflector, light is reflected about \mathbf{N} , so




- For a near-perfect reflector, you might expect the highlight to fall off quickly with increasing angle ω .
- Also known as:
 - “**rough specular**” reflection
 - “**directional diffuse**” reflection
 - “**glossy**” reflection

Specular Reflection Derivation



- One way to get this effect is to take $(\mathbf{R} \cdot \mathbf{V})$, raised to a power n_s
- As n_s gets larger,
 - the dropoff becomes {more,less} gradual
 - gives a {larger,smaller} highlight
 - simulates a {more,less} shiny surface

Specular Reflection



Iteration Three


- Since light is additive, we can handle multiple lights by taking the sum over every light.
- Our equation is now:

$$I = k_e + k_a I_a + \sum_i f(d_i) I_{li} \left[k_d (\mathbf{N} \cdot \mathbf{L}_i)_+ + k_s (\mathbf{V} \cdot \mathbf{R})_+^{n_s} \right]$$

- This is the **Phong illumination model**.
- Which quantities are spatial vectors?
- Which are RGB triples?
- Which are scalars?

Diffuse + Specular Reflection


- Diffuse + Specular



- You can control the ratio of diffuse to specular by adjusting k_s and k_d
- Contribution due to multiple light sources are simply added together

Specular Examples

- Effect on varying n_s




Choosing the Parameters?

- How would I model...
 - polished copper?
 - blue plastic?
 - lunar dust?


Choosing the Parameters

- N_s in the range $[0,100]$
- Try $K_a + K_d + K_s \leq 1$
- Use a small K_a (~ 0.1)

	N_s	K_d	K_s
Metal	Large	Small, color of metal	Large, color of metal
Plastic	Medium	Medium, color of plastic	Medium, white
Planet	0	Varying	0




**Appearance of a
diffuse (dull) sphere**




3D plot of reflected intensity

100% Diffuse




Appearance of a specular sphere





3D plot of reflected intensity

$$n = 15$$




$k_d = 0.7$
 $k_s = 0.4$
 $n = 15$






$$k_d = 0.85$$
$$k_s = 0.25$$
$$n = 15$$



Choosing the Parameters



Summary

- The most important thing to take away from this lecture is the final equation for the Phong model.
 - What is the physical meaning of each variable?
 - How are the terms computed?
 - What effect does each term contribute to the image?
 - What does varying the parameters do?